



## Revision of the genus *Suljuktocossus* Becker-Migdisova, 1949 (Hemiptera, Palaeontinidae), with description of a new species from Daohugou, Inner Mongolia, China

YING WANG & DONG REN<sup>1</sup>

College of Life Science, Capital Normal University, Beijing 100037, China

<sup>1</sup>Corresponding author. E-mail: rendong@mail.cnu.edu.cn

### Abstract

In this paper a complete specimen of fossil Palaeontinidae, *Suljuktocossus yinae* sp. nov. from Daohugou (Jiulongshan Formation) in Inner Mongolia, China is described. This new species is established based on both complete forewings and hindwings. According to this specimen, the diagnosis of the genus *Suljuktocossus* is revised. Moreover, based on the distribution of the genus *Suljuktocossus*, we consider the age of the Daohugou biota as Middle Jurassic.

**Key words** fossil, morphology, Cicadomorpha, Middle Jurassic

### Introduction

The genus *Suljuktocossus* erected by Becker-Migdisova is represented only by an incomplete forewing from the Early Jurassic of Shurab, Tadzhikistan (Becker-Migdisova, 1949). Recently another species belonging to this genus from the Daohugou biota, Inner Mongolia was reported (Wang *et al.*, 2007). From the same beds, the specimen we discovered is in remarkably good condition; both forewings and hind wings can be seen clearly.

Previously, the rich entomofauna from the Jiulongshan Formation in Daohugou Village, Ningcheng County, Inner Mongolia was dated as Middle Jurassic (Ren *et al.*, 1995; Ren & Krzemiski, 2002; Ren *et al.*, 2002; Shen *et al.*, 2003; Chen *et al.*, 2004; Liu *et al.*, 2004; Gao & Ren, 2006; Huang *et al.*, 2006; Ji *et al.*, 2006; Yao *et al.*, 2006; Tan & Ren, 2006; Liu *et al.*, 2007), Late Jurassic (Zhang, 2002) or Early Cretaceous (Wang *et al.*, 2005). Among these rich insect fossils, 14 genera and 27 species of palaeontinids were recorded from Daohugou biota (Wang *et al.*, 2006a-c; Wang *et al.*, 2007; Wang & Ren, 2006; Wang & Ren, 2007; Wang *et al.*, 2007a-c). This specimen of *Suljuktocossus* sheds new light on the age of the Daohugou fossil-bearing beds. Up to now, 3 species of *Suljuktocossus* are known from the Lower Jurassic of Shurab in Tadzhikistan and Daohugou, Inner Mongolia in China. These three species have a close relationship. Based on accurate Ar-Ar and SHRIMP U-Pb dating shows that the age of Daohugou intermediate-acid volcanic rocks overlying the Daohugou fossil-bearing beds is about 164–165 Ma, and that the age of this fossil-bearing beds is older than or equal to 165 Ma (Liu *et al.*, 2004). We adopt statement of Middle Jurassic age (Jiulongshan Formation) as proposed from the analysis of the Coleoptera (Tan & Ren, 2006), Plecoptera (Liu *et al.*, 2007) and Hemiptera fossils assemblage recently described by Yao *et al.* (2006) and Wang *et al.* (2006a).

## Material and methods

The specimen was examined by the dissecting microscope (Type: LEICA MZ12.5) and illustrated with the aid of a drawing tube. The specimen studied here is deposited in the Key Lab of Insect Evolution and Environmental Change, College of Life Science, Capital Normal University, Beijing, China. The wing venation nomenclature used in this paper is based on the interpretation of Becker-Migdisova (1949).

## Systematic descriptions

### Order Hemiptera Linnaeus, 1785

### Suborder Cicadomorpha Evans, 1946

### Family Palaeontinidae Handlirsch, 1906 =Cicadomorphidae Evans, 1956

### Genus *Suljuktocossus* Becker-Migdisova, 1949

Type species: *Suljuktocossus prosboloides* Becker-Migdisova, 1949.

**Revised diagnosis:** Forewing triangular with blurry CP and nodal line. Sc arising from basal single and parallel R before fusing with  $R_1$ . M 4-branched. CuA curved at basal, connected with M by a short crossvein m-cua, and dividing into  $CuA_1$  and  $CuA_2$ . Crossvein  $m_4$ -cua curved, and connected with CuA at the dividing point. CuP slightly curved before ending to margin. A 2-branched.  $A_2$  with 2 branches and the longer branch fused with  $A_1$  a little before distal end of clavus.

Hind wing with obvious indentation. Sc coalesced with R for a short distance and upwards single to indentation. M 4-branched.  $M_1$  separated from M basally, and fused with Rs for a long interval. The stem of  $M_{3+4}$  long and a little curved,  $M_{3+4}$  bifurcating in to  $M_3$  and  $M_4$  distad of CuA dividing point. CuA connected with  $M_{3+4}$  by a short crossvein. CuP simple. A with 2 branches.

**Species included:** Three species: *S. prosboloides*, Becker-Migdisova, 1949, *S. chifengensis* Wang, Zhang and Fang, 2007 and *S. yinae* sp. nov.

**Remarks:** Until now, 14 genera and 27 species assigned to the family Palaeontinidae have been reported from Daohugou, Inner Mongolia in China. The genus *Suljuktocossus* Becker-Migdisova, 1949 was erected just based on a single incomplete forewing. The material described here consists of complete forewings and hind wings with part and counterpart.

Among these genera, *Suljuktocossus* Becker-Migdisova, 1904 is similar to *Pseudocossus* Martynov, 1931 in forewing, but differs from the latter with blurry CP and nodal line, Sc without branches and the longer branch of  $A_2$  fused with  $A_1$  before distal end of clavus.


It differs from *Gansucossus* (Hong, 1983) Wang, Zhang and Fang, 2006 in hind wing with stem M long and  $A_1$  terminated in anal margin.

Based on hind wings venation (Sc coalesced with R for a short distance and upwards single to indentation,  $M_1$  fused with Rs for a long interval and  $M_{3+4}$  bifurcating in to  $M_3$  and  $M_4$  distad of CuA dividing point), this genus is quite similar to *Shurabocossus* Becker-Migdisova, 1949, but it differs from the latter mainly by the following combination of features: R 3-branched and  $A_2$  short and extended downwards to the anal margin near base.

### *Suljuktocossus yinae* sp. nov.

(Figs. 1, 2)

**Diagnosis:** Forewing with  $M_{1+2}$  separated distad of  $M_{3+4}$ ,  $m_4$ -cua slightly S-shaped and m-cua connected with M at the point of M arising from R+M. Hind wing as for genus.




**FIGURE 1.** *Suljuktocossus yinae* sp. nov., holotype: A, photograph of part, No. CNU-H-NN2007001-1; B, photograph of counterpart, No. CNU-H-NN2007001-2.

**Description.** Body compressed with both forewings and hind wings. Thorax and abdomen preserved, but obscure. One leg, probably hind leg, partly preserved. 2 segments of tarsi and second longer than first. Pretarsus visible.

Forewing length 56mm, width 27mm. Anterior margin blurry indented where nodal line reaches margin. CP present and ending at nodal indentation. Sc without branches, paralleling R+M for long interval and curved slightly upwards slightly before point of R+M separation, fused with  $R_1$  beyond point of R division into  $R_1$  and Rs. R+M arising basally. R bifurcating at same level of M separation into  $M_{1+2}$  and  $M_{3+4}$ .  $M_{1+2}$  separated

distad of  $M_{3+4}$  and divided into  $M_1$  and  $M_2$  slightly before indentation.  $M_1$  connected with Rs by short crossvein r-m. Crossvein m-cua connected with M at point of M separation from R+M. Stem of Cu short at base, forking into CuA and CuP. CuA curved when meeting crossvein m-cua, and bifurcating into  $CuA_1$  and  $CuA_2$  distad of M separated into  $M_{1+2}$  and  $M_{3+4}$ , but basas of  $M_{3+4}$  bifurcation. Crossvein  $m_4$ -cua slightly S-shaped and connected with CuA at point of CuA forking. CuP departing from Cu basally and slightly curved before reaching margin. Clavus small with 2 anal veins.  $A_2$  with 2 branches, shorter branch curved downward to margin at base; longer branch fused with  $A_1$  little before distal end of clavus. Nodal line traceable as indentation fused with  $R_1$  for a short distance and cutting Rs little beyond point of Sc fusing with  $R_1$ , then reaching M just beyond M fork, finally fused with  $m_4$ -cua to connection with  $CuA_2$ , following latter for a short distance and then reaching CuP at distal end of clavus.



**FIGURE 2.** *Suljuktocossus yinae* sp. nov., line drawing of holotype, No. CNU-H-NN2007001-1. *ts.* tarsus; *ptar*, pretarsus.

Hind-wing length 35mm, width 25mm. Hind-wing with obvious indentation. Sc arising basally, fused with R at the point of R branching into  $R_{1+2}$  and Rs, and extended a short distance, curved upwards single to indentation. R parallel Sc before branching into  $R_{1+2}$  and Rs.  $R_1$  arising from  $R_{1+2}$  at same level of indentation, then subparallel to Sc before ending at costal margin.  $R_2$  flattened before reaching costal margin above apex of wing. Rs curved and fused with  $M_1$  for a long interval after dividing from R. M 4-branched.  $M_1$  separated from M basally, and slightly convex before fusing with Rs.  $M_2$  straight and arising from M after the point of  $M_1$  separated from M. Stem of  $M_{3+4}$  long, connected with CuA by short crossvein  $m_{3+4}$ -cua and little curved when crossvein meeting  $M_{3+4}$ .  $M_{3+4}$  branched into  $M_3$  and  $M_4$  at same level of indentation. CuA bifurcated slightly basad than  $M_{3+4}$ . CuP simple. A 2-branched.  $A_2$  short and curved downwards to anal margin near base. Both forewings and hind wings with distinct color pattern consisting of regular dark patches on a pale ground.

**Etymology:** This species is named after Miss. Yin Ruiyue who collected this fossil and donated it to us.



**Material:** The new species is established based on compressed body with a pair of forewings and hindwings. Holotype: Both forewings and hindwings, part and counterpart, No.: CNU-H-NN2007001-1, CNU-H-NN2007001-2.

**Horizon and locality:** Jiulongshan Formation, Middle Jurassic, Daohugou Village, Ningcheng County, Inner Mongolia in China.

**Remarks:** This species is assigned to *Suljuktocossus* by the main characters of forewing, such as CP present, Sc arises from basal single and parallel R before fusing with R<sub>1</sub>, crossvein r-m short, the nodal line developed, are consistent with the type species.

The new species can be distinguished from the type species in the following features on the forewing: a short crossvein r-m near indentation, m<sub>4</sub>-cua slightly S-shaped and m-cua connected with M at the point of M arising from R+M.

This species is also similar to *S. chifengensis* Wang, Zhang and Fang, 2007, but it is different from the latter in the following features on the forewing: M<sub>1+2</sub> forking near indentation and M<sub>4</sub> not sharply curved.

## Acknowledgements

We sincerely thank Dr. Shih ChungKun (College of Life Sciences, Capital Normal University) for his improvement of our manuscript. This research was supported by grants from the National Nature Science Foundation of China (No. 30430100) and the PHR (IHLB) Project of Beijing Municipal Commission of Education.

## References

- Becker-Migdisova, E.E. (1949) Mesozoic Homoptera of Middle Asia. *Trudy Paleontologicheskogo Instituta AN SSSR*, 22, 1–68 (in Russian).
- Chen, W., Ji, Q., Liu, D.Y., Zhang, Y., Song, B. & Liu, X.Y. (2004) Isotope geochronology of the fossil-bearing beds in the Daohugou area, Ningcheng, Inner Mongolia. *Geological Bulletin of China*, 23(12), 1165–1169 (in Chinese with English abstract).
- Gao, K.Q. & Ren, D. (2006) Radiometric dating of ignimbrite from Inner Mongolia provides no indication of a post-Middle Jurassic age for the Daohugou beds. *Acta Geologica Sinica*, 80, 42–45.
- Huang, D.Y., Nel, A., Shen, Y.B., Selden P. A. & Lin, Q.B. (2006) Discussions on the age of the Daohugou faunaevidence from invertebrates. *Progress in Natural Science*, 16, 308–312.
- Ji, Q., Luo, Z.X., Yuan, C.X. & Tabrum, A.R. (2006) A swimming mammaliaform from the Middle Jurassic and ecomorphological diversification of early mammals. *Science*, 311, 1123–1127.
- Liu, Y.Q., Liu, Y.X., Li, P.X., Zhang, H., Zhang, L.J., Li, Y. & Xia, H.D. (2004) Daohugou biota-bearing lithostratigraphic succession on the southeastern margin of the Ningcheng basin, Inner Mongolia, and its geochronology. *Geological Bulletin of China*, 23(12), 1180–1185 (in Chinese with English abstract).
- Liu, Y.S., Ren, D., Sinitshenkova, N. & Shih, C.K. (2007) The oldest known record of Taeniopterygidae in the Middle Jurassic of Daohugou, Inner Mongolia, China (Insecta: Plecoptera). *Zootaxa*, 1521, 1–8.
- Ren, D., Lu, L.W., Guo, Z.G. & Ji, S.A. (1995) *Faunae and Stratigraphy of Jurassic-Cretaceous in Beijing and the Adjacent Areas*. Seismic Publishing House, Beijing, pp. 64–73.
- Ren, D., Yin, J.C. & Dou, W.X. (1998) Late Jurassic palaeontinids Homoptera: Auchenorrhyncha from Heibei and Liaoning Provinces in China. *Entomologia Sinica*, 5(3), 222–232.
- Ren, D. & Krzemiski, M. (2002) Eoptychopterridae (Diptera) from the Middle Jurassic of China. *Annales Zoologici*, 52(2), 207–210.
- Ren, D., Gao, K.Q., Guo, Z.G., Ji, S.A., Tan, J.J. & Song, Z. (2002) Stratigraphic division of the Jurassic in the Daohugou area Ningcheng, Inner Mongolia. *Geological Bulletin of China*, 21(8-9), 584–591 (in Chinese with English abstract).
- Shen, Y.B., Chen, P.J. & Huang, D.Y. (2003) Age of the fossil conchostracans from Daohugou of Ningcheng, Inner Mongolia. *Journal of Stratigraphy*, 27(4), 311–313 (in Chinese with English abstract).
- Tan, J.J. & Ren, D. (2006) Jurassic and Cretaceous Cupedomorpha (Insecta: Coleoptera) faunas of China. *Progress in Natural Science*, 16(Special Issue), 313–322.

- Wang, B., Zhang, H.C. & Fang, Y. (2006a) Some Jurassic Palaeontinidae (Insecta, Hemiptera) from Daohugou, Inner Mongolia, China. *Palaeoworld*, 15, 115–125.
- Wang, B., Zhang, H.C. & Fang, Y. (2006b) *Gansucossus*, a replacement name for *Yumenia* Hong, 1982 (Insecta, Hemiptera, Palaeontinidae), with description of a new genus. *Zootaxa*, 1268, 59–68.
- Wang, B., Zhang, H.C. & Fang, Y. & Duan, Y. (2006c) Revision of the genus *Sinopalaeocossus* Hong (Hemiptera: Palaeontinidae), with description of a new species from the Middle Jurassic of China. *Zootaxa*, 1349, 37–45.
- Wang, B., Zhang, H.C. & Fang, Y. (2007) Middle Jurassic Palaeontinidae (Insect, Hemiptera) from Daohugou of China. *Alavesia*, 1, 89–104.
- Wang, X.L., Zhou, Z.H., He H.Y., Jin, F., Wang, Y.Q., Zhang, J.Y., Wang, Y., Xu, X. & Zhang, F.C. (2005) Stratigraphic relation and age discussion of the Daohugou area, Ningcheng, Inner Mongolia. *Chinese Science Bulletin*, 50(19), 2127–2135.
- Wang, Y. & Ren, D. (2006) Middle Jurassic *Pseudocossus* fossils from Daohugou, Inner Mongolia in China (Homoptera, Palaeontinidae). *Acta Zootaxonomica Sinica*, 31(2), 289–293 (in Chinese with English abstract).
- Wang, Y. & Ren, D. (2007) Two new genera of fossil palaeontinids from the Middle Jurassic in Daohugou, Inner Mongolia, China (Hemiptera, Palaeontinidae). *Zootaxa*, 1390, 41–49.
- Wang, Y., Ren, D. & Shih, C.K. (2007a) Discovery of Middle Jurassic palaeontinids from Inner Mongolia, China (Homoptera: Palaeontinidae). *Progress in Natural Science*, 17(1), 112–116.
- Wang, Y., Ren, D. & Shih, C.K. (2007b) New discovery of palaeontinid fossils from the Middle Jurassic in Daohugou, Inner Mongolia (Homoptera, Palaeontinidae). *Science in China (Ser. D Earth Sciences)*, 50(4), 481–486.
- Yao, Y.Z., Cai, W.Z. & Ren, D. (2006) The first discovery of fossil rhopalids (Heteroptera: Coreoidea) from Middle Jurassic of Inner Mongolia, China. *Zootaxa*, 1269, 57–68.
- Zhang, J.F. (2002) Discovery of Daohugou biota (pre-Jehol biota) with a discussion on its geological age. *Journal of Stratigraphy*, 26(3), 173–177, 215 (in Chinese with English abstract).